

Eyes Over Puget Sound: Strategies to increase the impact and effectiveness of long-term marine PERTITENT OF ECOLOGY State of Washington





Measuring long-term trends in eutrophication, dissolved oxygen and physical variables



Water Quality variables measured monthly at 27 stations




Physical variables

- Temperature
- Salinity
- Density

Chemical variables

- Oxygen
- Nitrate
- Silicate
- Phosphate
- Ammonium
- Nutrient ratios
- pH





Bio-optical variables

- Water clarity
- · Chlorophyll a
- Euphotic depth

Monthly Baselines 1999-2008

Strength of long-term monitoring programs

- Different perspective on processes operating on larger temporal and spatial scales
- 2. Opportunity for understanding variability and influencing factors over different scales
- 3. Objectivity through data
- 4. Warning systems for environmental change
- 5. Temporal and spatial context supporting focused studies
- 6. Logistical infrastructure and expertise

Despite its undisputed value as a quantitative instrument, maintaining and committing to long-term monitoring programs is challenging:

- 1. Resource competition with short-term studies
- 2. Inconsistency in funding and staff
- 3. Changes in methods as technology evolves
- 4. Lack of standardized procedures and consistent data quality
- 5. Wide range of temporal and spatial scales
- 6. Trade-offs between representative measurements and local impacts

Dilemma of long-term monitoring programs providing exclusively data

•Data collection, management, QC, and instrument maintenance often dominate the activities to the detriment of data workup and presentation.

•Providing only data limits the program to shine to a hand full of data users. → Few people that can advocate for the value of the program.

The combination of data and stories let a program shine!

How do you create that without additional resources?

Strategies to increase the impact and effectiveness of long-term marine monitoring programs (story material)

We humans like to think on human scales (time and space)

Image rich to engage more people



Headline driven information bits

- •The ocean and air remain warm with sunshine and dry weather across the region.
- •Puget Sound is a lot warmer going into the new year.
- •Hood Canal is, breaking its low temperature stint.
- •Patches of jellyfish are overwintering in finger inlets of South Sound.

Deliver on a time scale that humans like to think in (within a few days)

Public **Progress** Interest Relevant Persistent Documentation Visible Organized **Program** Efficient Funded Engaging Valuable Progressive Scale Effective Variables Resolution Representative Accurate/Precise Innovative Continuous **Back compatibility** Collaborative Story driven Free Knowledgeable Motivated Invested Informative Accessible **Evaluation** Understandable **Awareness** Engagement

What is EOPS?

Just collecting data and writing reports is not enough...



Eyes Over Puget Sound





EOPS

Multiple perspectives...



Be informed within 2 days



- 1. Everybody is informed about their backyard marine environment.
- Free educational material.
- 3. Access for more information + data.

EOPS offers multiple perspectives

Different levels of detail (overview -> detail)

 Present observations in historical context (temporal context)

 Observations from different vantage points (spatial context)

Beauty and challenges of Puget Sound




EOPS high public resonance

Steady interest

> 600 Listserve subscribers

General interest

Over 2 million downloads!






Who are the potential users?

Listserver 600 subscribers

Avg 46,000 downloads per edition

Public readers not accounted for

EOPS spreads via social media



What does EOPS provide?

TEMPORAL PERSPECTIVE:

A combined presentation of present field conditions and 14 years of historical data.

SPATIAL PERSPECTIVE:

A spatial and temporal coverage that matches the scale of local, climatic and oceanic processes that permeate the monitoring network.

BEING INFORMED ABOUT THE PRESENT CONDITION, (useful for decision making)

Only information, image-rich-story-driven, published two days after visiting the field.

MORE INVESTED AND EMPOWERED PUBLIC:

EOPS creates impact on a very tangible, human scale. Staff gain visibility and positive feedback and are also more invested.

IMPROVED WORKFLOW AND BETTER DATA

Public visibility motivates!

Monthly reporting with a 2-day turnaround streamlines operations.


How do you do that without additional resources?

The benefits of combining QC routines with environmental condition updates

- QC in temporal, spatial context results in better data quality
- QC on present/recent conditions:
 - reduce error propagation
 - prevent backlogs
 - establish sense of control and relevance
 - fosters positive feedback
- Reporting relative to a baselines facilitates stories telling, better QC and easier communication across programs.

A focus on anomalies combines temporal scales and <u>leverages</u> the strength of long-term monitoring programs

Ocean boundary conditions impact PS water quality



n.a.

n.a.

How do we do it?

Leveraging our flight time

- Empty transit flights
 Seattle Olympia with
 camera on board
- Document blooms, debris, animal aggregations, oil sheens, water boundaries
- Unique perspective, minimal extra cost





Eyes Over Puget Sound





Marine conditions from 2-17-2015 at a glance



Field log

LONG-TERM

Climate

Water column

Aerial photos

Ferry and Satellite

Moorings

Mya Keyzers Laura Hermanson Joe Leatherman





Skip Albertson



Julia Bos Suzan Pool



Dr. Christopher Krembs



Dr. Brandon Sackmann,

Please give us feedback

Personal field log

p. 4

Why do I love Puget Sound?

Climate conditions

The air temperatures and ocean conditions remain warm. Rivers are above normal except at the coast (Chehalis). The past week has generally been sunny and dry across the region.

Water column

Puget Sound is a lot warmer at the end of 2014 with new maximum temperatures observed throughout the Sound! The higher dissolved oxygen and cold temperature anomalies in Hood Canal are disappearing.

Moorings

p. 37

Compared to past years, 2015 starts with warmer water and lower salinity at Mukilteo. Dissolved oxygen is rebounding.

Aerial photography

p. 10

Patches of jellyfish persist in finger inlets of South Sound. First signs of growing phytoplankton are seen where water is turning green. Otherwise, the surface waters are nice shades of blue and green and, as expected, carry sediment near rivers.

Ferry and satellite

Victoria Clipper is in the shipyard

p. 36

Editorial assistance provided by:

Suzan Pool, Carol Maloy



Personal Field Impression 2-15-15



Field log

Climate

Water column

Aerial photos

Ferry and Satellite

Moorings



What do you love the most about Puget Sound?









Wondrous marine life

Islands and hidden bays

Charismatic marine mammals



Personal Field Impression 2-15-15



Field log

Climate

Water column

Aerial photos

Ferry and Satellite

Moorings



Miles of diverse coastline





Glacial flour and river plumes



Unique geological features



Marine conditions from 2-17-2015 at a glance



Field log

LONG-TERM

Climate

Water column

Aerial photos

Ferry and Satellite

Moorings

Mya Keyzers Laura Hermanson Joe Leatherman



Skip Albertson



Julia Bos Suzan Pool



Dr. Christopher Krembs



Guest: Dr. Brandon Sackmann, Integral Please give us feedback

Personal field log Why do I love Puget Sound?

<u>p. 4</u>

Climate conditions

p. 6

The air temperatures and ocean conditions remain warm. Rivers are above normal except at the coast (Chehalis). The past week has generally been sunny and dry across the region.

Water column

<u>p. 7</u>

Puget Sound is a lot warmer at the end of 2014 with new maximum temperatures observed throughout the Sound! The higher dissolved oxygen and cold temperature anomalies in Hood Canal are disappearing.

Moorings

<u>p. 37</u>

Compared to past years, 2015 starts with warmer water and lower salinity at Mukilteo. Dissolved oxygen is rebounding.

Aerial photography

p. 10

Patches of jellyfish persist in finger inlets of South Sound. First signs of growing phytoplankton are seen where water is turning green. Otherwise, the surface waters are nice shades of blue and green and, as expected, carry sediment near rivers.

Ferry and satellite

Victoria Clipper is in the shipyard

<u>p. 36</u>

Editorial assistance provided by:

Suzan Pool, Carol Maloy



Climate and natural influences before 2-17-2015



Field log

Climate

Water column

Aerial photos

Ferry and Satellite

Moorings



New section! Climate and natural influences are conditions that influence our marine waters, including weather, rivers, and the adjacent ocean (previously called Weather). For an explanation of the figure, see: http://www.ecy.wa.gov/programs/eap/mar_wat/weather.html, page 26.

Summary:


Air temperatures remain above normal, continuing a 10-month trend.

Precipitation has been below normal for the past week and month in the Central Puget Sound region.

Sunshine has been abundant for the past five days and above normal for the winter.

River flows are above normal across the Puget Sound region, but below normal at the coast (Chehalis River).

PDO remains in the warm phase, and upwelling is below normal.





Marine conditions from 2-17-2015 at a glance



Field log

LONG-TERM

Climate

Water column

Aerial photos

Ferry and Satellite

Moorings

Mya Keyzers Laura Hermanson Joe Leatherman





Skip Albertson



Julia Bos Suzan Pool



Dr. Christopher Krembs



Dr. Brandon Sackmann,

Please give us feedback

Personal field log Why do I love Puget Sound?

p. 4

Climate conditions

The air temperatures and ocean conditions remain warm. Rivers are above normal except at the coast (Chehalis). The past week has generally been sunny and dry across the region.

Water column

Puget Sound is a lot warmer at the end of 2014 with new maximum temperatures observed throughout the Sound! The higher dissolved oxygen and cold temperature anomalies in Hood Canal are disappearing.

Moorings

p. 37

Compared to past years, 2015 starts with warmer water and lower salinity at Mukilteo. Dissolved oxygen is rebounding.

Aerial photography

p. 10

Patches of jellyfish persist in finger inlets of South Sound. First signs of growing phytoplankton are seen where water is turning green. Otherwise, the surface waters are nice shades of blue and green and, as expected, carry sediment near rivers.

Ferry and satellite

Victoria Clipper is in the shipyard

p. 36

Editorial assistance provided by:

Suzan Pool, Carol Maloy

Physical conditions tracked in statistically historic context



Field log

Weather Water column

Aerial photos

Ferry and Satellite


Moorings

In 2014, conditions were dominated by <u>warm water</u> during summer and fall, associated with the NE Pacific Ocean warm surface anomaly. In October and November, temperatures were the highest on our record since 1989. In 2014, salinity was higher and later, waters south of Admiralty Reach became fresher. Oxygen was mostly lower except in Hood Canal where a high anomaly persisted into the fall.

Nov. 2014:

Red boxes show that the water measured in fall 2014 is warmer than any of our measurements since 1999

Explore profiles



= higher than expected (>IQR, n=13)

higher than previous measurements

= expected (=IQR, n=13)

= no data

= lower than expected (>IQR, n=13)= lower than previous measurements



Marine conditions from 2-17-2015 at a glance



Field log

LONG-TERM

Climate

Water column

Aerial photos

Ferry and Satellite

Moorings

Mya Keyzers Laura Hermanson Joe Leatherman





Skip Albertson



Julia Bos Suzan Pool



Guest: Dr. Brandon Sackmann,

Dr. Christopher

Krembs

Integral



Please give us feedback

Personal field log Why do I love Puget Sound?

<u>p. 4</u>

Climate conditions

p. 6

The air temperatures and ocean conditions remain warm. Rivers are above normal except at the coast (Chehalis). The past week has generally been sunny and dry across the region.

Water column

p. 7

Puget Sound is a lot warmer at the end of 2014 with new maximum temperatures observed throughout the Sound! The higher dissolved oxygen and cold temperature anomalies in Hood Canal are disappearing.

Moorings

p. 37

Compared to past years, 2015 starts with warmer water and lower salinity at Mukilteo. Dissolved oxygen is rebounding.

Aerial photography

p. 10

Patches of jellyfish persist in finger inlets of South Sound. First signs of growing phytoplankton are seen where water is turning green. Otherwise, the surface waters are nice shades of blue and green and, as expected, carry sediment near rivers.

Ferry and satellite

Victoria Clipper is in the shipyard

p. 36

Editorial assistance provided by:

Suzan Pool, Carol Maloy








Aerial photography 1-15-2013










Aerial photography 1-15-2013








Aerial photography 12-13-2012 (1)






Willapa shellfish grower applying local ground up oyster shells to improve clam substrate. Location: West of Bone River (Willapa Bay), 12:53 PM





Aerial photography 11-08-2012






Red-brown algal bloom and jellyfish and turquoise water. Location: Eld Inlet (South Sound), 12:56 PM



Aerial photography









Aerial photography 2-26-2013



Navigate



Large tidal eddy carrying suspended sediment with debris lines.

Location: Nisqually Reach looking into Dana Passage (Central Sound), 3:08 PM







Navigate

Aerial photos Ferry and Satellite Field log Weather Water column Moorings Suspended sediment





Aerial photography 11-2-2010



Navigate


Aerial photos Ferry and Satellite Field log Weather Water column Moorings





Aerial photography 7-31-2012











Navigate



Long foam lines persist after ships have passed - Elliott Bay to Admiralty Reach, 3:38 PM


















Aerial photography 9-11-2012



Navigate

Aerial photos Field log Water column Ferry and Satellite Moorings Weather







Navigate




Red-brown bloom. Location: Quartermaster Harbor (Vashon Island), 4:32 PM



Aerial photography 7-31-2012









Aerial photography 7-31-2012



Navigate

Aerial photos Water column Ferry and Satellite Moorings Field log Weather ship tracks

Red-brown and turquoise blooms, jellyfish (+ cloud reflections). Location: Budd Inlet (South Sound), 4:56 PM

Microbial food web: The importance of the micro-grazer Noctiluca in Puget Sound



Noctiluca surface aggregations looking south into Central Basin, Puget Sound, Washington, USA






Aerial photography June 2011 📵



Navigate



Noctiluca filaments drifting at the surface east of Vashon Island, looking onto Des Moines. The eastern tip of Vashon Island can be seen to the right 4:44 PM





Aerial photography 6-12-2012










Aerial photography 6-12-2012







Aerial photography 9-11-2012









Aerial photography 2-26-2013







Ferry and Satellite

Brandon Sackmann

Ferry Monitoring



Water Quality Indicators

(Deployed 14 May 2010)

- Phytoplankton (Chlorophyll a fluor.)
- Turbidity
- Sea Surface Temperature
- River water (CDOM)
- Salinity (Apr. 2012)







Ferry Monitoring




- Advantages
 - Cost-effective data collection
 - 100 m spatial resolution (5 sec.)
 - 4-hr temporal resolution
 - Regular schedules/Reliable
 - 80 mile long transect (35 mph)
 - 1-2 times daily (year-round)
- Daily data pickup
- Simplicity helps ensure a sustainable program...

Seaplane → Ferry → Satellite



Chlorophyll












Field log Climate Water column Aerial photos Ferry and Satellite Moorings





Current Conditions:


Bloom in central Puget Sound begins to fade as temperatures cool; max temperatures generally <15 °C. MODIS reveals extensive bloom at entrance to Strait of Juan de Fuca. Thermal imagery from Landsat 8 shows warmer water in Strait of Georgia, Whidbey Basin, and finger inlets of South Puget Sound.
















- **B.** Stratification shows signs of weakening (temperature is declining indicating mixing) and bloom is dissipating.
- **C.** Weak tides in August and September associated with warmer temperatures in Strait of Juan de Fuca.





Victoria Clipper rendezvous near Kingston













Field log Climate Water column Aerial photos Ferry and Satellite Moorings



MODIS-Terra (top) and MODIS-Aqua (bottom) reveal intense offshore bloom near entrance to Strait of Juan de Fuca!



Imagery obtained from NASA's OceanColor WEB


http://oceancolor.gsfc.nasa.gov/








Field log Climate Water column Aerial photos Ferry and Satellite Moorings



Thermal imagery from the Landsat 8 satellite show warm waters throughout Strait of Georgia and Whidbey Basin (left). Warmer temperatures were in finger inlets throughout South Puget Sound; cooler temperatures highlight areas experiencing increased mixing (top).

Hypothesis for combining a series of recent observations affecting energy and material transfer to higher trophic levels



Hypothesis!

Increases in nitrate concentrations could be caused by a top-down control on phytoplankton biomass.

Is Noctiluca
a visible
harbinger of a
food web
change?

Follow the experts
WebEx

Are changes in higher tropic levels part of a story of the low food web?



EOPS



Increased focus on the environment

...Others are more engaged

We do a better job...

Information is better

